



Jurnal Kedokteran dan Kesehatan Indonesia

Indonesian Journal of Medicine and Health

Journal homepage : www.journal.uui.ac.id/index.php/jkki

Alzheimer disease and acetylcholinesterase inhibitors

Zahia Kabouche¹

¹Université des frères Mentouri-Constantine, Département de chimie, Laboratoire d'Obtention de Substances Thérapeutiques (LOST), 25000 Constantine (Algeria)

EDITORIAL

ARTICLE INFO

*Corresponding author:
zahiakabouche@umc.edu.dz

DOI : 10.20885/JKKI.Vol7.Iss5.art1

Alzheimer's disease (AD) or Alzheimer's is a chronic neurodegenerative disorder. It's characterized by loss of neurons and synapses in the cerebral cortex and certain subcortical regions. AD patients are identified by problems with memory and language, getting lost and desorientation. AD affects about 6% of people 65 years and older but 4% to 5% of cases are younger.¹ It is one of the most costly diseases in the world. In 2015, there were about 48 million people worldwide with AD.

To prevent AD, intellectual activities such as reading and playing chess are recommended. In muslim countries, learning quran is linked to a reduced risk of AD. The cause for most Alzheimer's cases is still mostly unknown except for 1% to 5% of cases linked to genetic factors.

AD has been identified as a protein misfolding disease (proteopathy), caused by plaque accumulation of abnormally folded amyloid beta protein in the brain.² However, there is another hypothesis which proposes that AD is caused by reduced synthesis of the neurotransmitter acetylcholine.³ The enzyme Acetylcholinesterase breaks acetylcholine down after acetylcholine is used in the brain.

Galanthamine, a natural product isolated from *Galanthus nivalis*, *G. woronowii* and some other Amaryllidaceae plants and rivastigmine which is a natural derived compound and Donepezil which is a synthetic compound are used as cholinesterase inhibitors.

Polyphenols which have been shown to possess antioxidant activity are the principal constituents of the mediterranean diet which is known to decrease the risk of AD. Antioxidants may slow the progression of AD and minimise neuronal degeneration.⁴ The compounds that exhibit anticholinesterase activity are also related to radical scavenging activity.⁵ Polyphenols and terpenoids from Lamiaceae plants possess antioxidant and anticholinesterase activities.^{6,7}

REFERENCES:

- Centers for Disease Control and Prevention (CDC). Zika virus [Internet]. 2016. Available from: <http://www.cdc.gov/zika/index.html>
- Kutsuna S, Kato Y, Takasaki T, Moi ML, Kotaki A, Uemura H, et al. Two cases of Zika fever imported from French Polynesia to Japan, December 2013 to January 2014. *Euro Surveillance*. 2014;19(4):50-3.
- Petersen LR, Jamieson DJ, Powers AM, Honein MA. Zika Virus. *New England Journal of Medicine*. 2016;374(16):1552-63.

4. Karim O, Goorhuis A, Schinkel J, Codrington J, Vreden SGS, Vermaat JS, et al. Thrombocytopenia and subcutaneous bleedings in a patient with Zika virus infection. *Lancet*. 2016;387(10022):939–40.
5. Rozé B, Najioullah F, Fergé J, Apetse K, Brouste Y, Cesaire R, et al. Zika virus detection in urine from patients with Guillain-Barré syndrome on Martinique, January 2016. *Euro Surveill*. 2016 Jan;21(9):30154.
6. Lanciotti RS, Kosoy OL, Laven JJ, Velez JO, Lambert AJ, Johnson AJ, et al. Genetic and serologic properties of Zika virus associated with an epidemic, Yap State, Micronesia, 2007. *Emerging Infectious Diseases*. 2008;14(8):1232–9.
7. Yilmaz A, Boğa M, Topçu G. Novel terpenoids with potential Anti-alzheimer activity from *Nepeta obtusifolia*. *Records of Natural Products*. 2016;10(5):530–41.